

Wolfgang Husinsky

TU Wien, Institut für Angewandte Physik

Tuesday, 15th January 2019, 16:00 s.t.

TU Wien, Institut für Angewandte Physik, E134

1040 Wien, Wiedner Hauptstraße 8-10

Yellow Tower „B“, 5th floor, SEM.R. DB gelb 05 B



Experiments, Finite-Element and Molecular Dynamic Calculations for LIPSS (Laser Induced Periodic Surface Structures) formation during ultra-short laser ablation

Irradiation of many surface materials by ultra-short laser radiation, typically with pulsewidths of 10-300fs, results in the development of Laser-induced self-organized periodic surface structures (ripples). The periodicity of these structures range from the wavelength λ of the laser to well below λ . Structures down to a few tens of nm can be observed. However, the existence and form of the LiPSS critically depends on the the intensity, fluence, wavelength and pulsewidth of the laser light as well as on the number of laser pulses and, not to forget, the material properties [1].

The development of the LIPSS structure can be regarded as a complex interplay between the electric field on the surfaces (including surface plasmon-polaritons) [2] and the ultra-short-laser-ablation dynamics [3]. Since the formation of LIPSS in most cases requires multiple laser pulses, the surface topography and hence the electric field distribution will change from shot to shot. Only the resulting feedback allows for the development of the LIPSS.


The seminar discusses a new approach to simulate the physics described above by calculating the electric field on and below the surface after each laser pulse for the newly developed surface topology with COMSOL Multiphysics (finite element method). This information of this field distribution, which should incorporate all possible plasmons etc., is then transferred to the irradiated material (various possibilities according to the wealth of physical processes in laser ablation [3] are evaluated) as a starting point for Molecular Dynamic (MD) simulations (LAMMPS) in order to obtain the new surface topology.

References

[1] Nathala, C. S. R., A. Ajami, A. A. Ionin, S. I. Kudryashov, S. V. Makarov, T. Ganz, A. Assion and W. Husinsky, *Optics Express* **23(5)**, 5915 (2015).

[2] Kudryashov, S. I., S. V. Makarov, A. A. Ionin, C. S. R. Nathala, A. Ajami, T. Ganz, A. Assion and W. Husinsky, *Optics Letters* **40(21)**, 4967 (2015).

[3] Dachraoui, H. and W. Husinsky., *Applied Physics Letters* **89(10)**, 4102 (2006).



Result from a COMSOL simulation of the square of the electric field for a $2\mu\text{m}$ by $2\mu\text{m}$ Ti surface with a thin oxide over-layer irradiated with a constant laser intensity.

All interested colleagues are welcome to this seminar lecture (45 min. presentation followed by discussion).

Friedrich Aumayr
(LVA-Leiter)

Richard Wilhelm
(Seminar Chair)